

## Research

# Lipid functions in cytochrome *bc* complexes: an odd evolutionary transition in a membrane protein structure

S. Saif Hasan and William A. Cramer\*

Department of Biological Sciences, Purdue University, West Lafayette, IN 47907, USA

Lipid-binding sites and properties were compared in the hetero-oligomeric cytochrome (cyt) *b<sub>6</sub>f* and the yeast *bc<sub>1</sub>* complexes that function, respectively, in photosynthetic and respiratory electron transport. Seven lipid-binding sites in the monomeric unit of the dimeric cyanobacterial *b<sub>6</sub>f* complex overlap four sites in the *Chlamydomonas reinhardtii* algal *b<sub>6</sub>f* complex and four in the yeast *bc<sub>1</sub>* complex. The proposed lipid functions include: (i) interfacial–interhelix mediation between (a) the two 8-subunit monomers of the dimeric complex, (b) between the core domain (cyt *b*, subunit IV) and the six trans membrane helices of the peripheral domain (cyt *f*, iron–sulphur protein (ISP), and four small subunits in the boundary ‘picket fence’); (ii) stabilization of the ISP domain-swapped trans-membrane helix; (iii) neutralization of basic residues in the single helix of cyt *f* and of the ISP; (iv) a ‘latch’ to photosystem I provided by the β-carotene chain protruding through the ‘picket fence’; (v) presence of a lipid and chlorophyll *a* chlorin ring in *b<sub>6</sub>f* in place of the eighth helix in the *bc<sub>1</sub>* cyt *b* polypeptide. The question is posed of the function of the lipid substitution in relation to the evolutionary change between the eight and seven helix structures of the cyt *b* polypeptide. On the basis of the known n-side activation of light harvesting complex II (LHCII) kinase by the p-side level of plastoquinol, one possibility is that the change was directed by the selective advantage of p- to n-side trans membrane signalling functions in *b<sub>6</sub>f*, with the lipid either mediating this function or substituting for the trans membrane helix of a signalling protein lost in crystallization.

**Keywords:** assembly; cytochrome complex; LHCII kinase; quinol/quinone; super-complex; trans membrane signalling

## 1. INTRODUCTION

The crystal structure of the 230 kDa (including prosthetic groups) symmetrically dimeric cytochrome (cyt) *b<sub>6</sub>f* complex of oxygenic photosynthesis has been determined for the filamentous cyanobacteria *Mastigocladus laminosus* [1–4] and *Nostoc* PCC 7120 [5], and the green alga *Chlamydomonas reinhardtii* [6]. The Protein Data Bank (PDB) accession numbers are, respectively, (i) *M. laminosus*: 2E74 (native), 2E75 (with bound electrochemically negative (n)-side inhibitor, 2-nonyl-4-hydroxyquinoline N-oxide, NQNO), 1VF5 and 2E76 (with tridecyl-stigmatellin, usually considered an electrochemically positive (p)-side inhibitor, but found at both n- and p-side quinone-binding sites), 2D2C (with p-side inhibitor, DBMIB); (ii) *Nostoc*: 2ZT9 (native); (iii) *C. reinhardtii*: 1Q90 (with tridecyl-stigmatellin). The crystal structures are consistent in describing eight polypeptide subunits, 13 trans membrane helices (TMHs) per monomer and seven prosthetic groups (four haems (haem *f* on the p-side luminal phase, two *b*-type haems, *b<sub>p</sub>* and *b<sub>n</sub>*, on the

p- and n-sides of the complex, covalently bound haem *c<sub>n</sub>* whose central Fe atom is within 4 Å of the propionate of haem *b<sub>n</sub>*), one [2Fe–2S] cluster at the p-side interface, one chlorophyll *a*, and one β-carotene]. Four of the five redox prosthetic groups, with the exception of haem *c<sub>n</sub>*, have the corresponding functional groups haems *c<sub>1</sub>*, *b<sub>p</sub>* and *b<sub>n</sub>*, and the [2Fe–2S] cluster, in the cyt *bc<sub>1</sub>* complex.

The core of the cyt *b<sub>6</sub>f* complex consists of: (i) the four TMHs of the cyt *b* subunit containing two trans membrane haems, which is structurally and functionally equivalent to the N-terminal trans membrane helical (helices A–D) haem-binding domain of the eight TMH cyt *b* subunits in the *bc<sub>1</sub>* complex (reviewed recently in [7]); (ii) subunit IV, with three TMHs that are structurally homologous to TMHs E–G of the *bc<sub>1</sub>* complex. The peripheral domain of the *b<sub>6</sub>f* complex consists of six single TMH subunits, cyt *f* [8], the Rieske iron–sulphur protein (ISP) [9] and the small photosynthetic electron transport (Pet) subunits G, N, L and M that have been described as ‘hydrophobic sticks’ [10]. A ribbon diagram of the structure derived from PDB 2E74 is shown in figure 1*a*.

A dependence of the membrane protein crystal structures on the presence of specific lipids has been demonstrated for a number of membrane proteins and protein complexes. These include the photosynthetic light-harvesting chlorophyll protein [11], the bacterial

\* Author for correspondence (waclab@purdue.edu).

One contribution of 16 to a Theo Murphy Meeting Issue ‘The plant thylakoid membrane: structure, organization, assembly and dynamic response to the environment’.




Figure 1. Cytochrome *b<sub>6</sub>f* complex of *Mastigocladus laminosus* (PDB 2E74). (a) Redox groups in the cyt *b<sub>6</sub>f* complex. The transmembrane *b*-haems (*b<sub>p</sub>* and *b<sub>n</sub>*) are shown in green, and haem *b<sub>c</sub>* in black. Haem of cyt *f* soluble domain is shown in red; [2Fe-2S] cluster of the ISP is shown as brown and yellow spheres (brown, Fe; yellow, S). The eight polypeptides in cyt *b<sub>6</sub>f* monomer are shown as ribbons (cyt *f*, cyan; cyt *b<sub>6</sub>*, light brown; ISP, pink; subIV, yellow; Pet G, gray; Pet L, green; Pet M, wheat; Pet N, dark blue). (b) Lipid-, detergent- and pigment-binding sites in cyt *b<sub>6</sub>f*. Natural acidic sulfolipid is shown in wheat. Natural galactolipid, monogalactosyl-diacylglycerol (MGDG), in *C. reinhardtii* *b<sub>6</sub>f* structure (PDB 1Q90) are in cyan and red. The synthetic lipid, dioleoylphosphatidylcholine (DOPC), used for crystallization of *b<sub>6</sub>f* from *M. laminosus* and *Nostoc* PCC 7120 is shown in red. Ordered molecules of the detergent *n*-undecyl-β-D-maltopyranoside (UDM) found in the cyanobacterial *b<sub>6</sub>f* structures are shown as, white and red sticks. Native chlorophyll *a* (chl-*a*), β-carotene (β-car) and eicosane (from *C. reinhardtii* *b<sub>6</sub>f*) are shown as green, yellow and dark blue sticks. Figure was generated by superposition of *C. reinhardtii* (PDB 1Q90) and *M. laminosus* *b<sub>6</sub>f* (PDB 1VF5, 2E74).

photosynthetic reaction centre [12–16], photosynthetic reaction centres II [17–19] and I [18,20], bovine [21] and bacterial cyt oxidase [22,23] and the cyt *bc<sub>1</sub>* complex [24].

## 2. DISCUSSION

Lipid functions in the *b<sub>6</sub>f* complex, described in figure 1*b*, include [25]: first, n-side and p-side stabilization of the dimer structure, the former analogous to function of the second cardiolipin in the *bc<sub>1</sub>* complex (yeast; 3CX5 [26]); and secondly, n-side stabilization of the domain-swapped trans membrane helices of the high potential Rieske ISP by anionic sulpholipid [25,27], and by phosphatidic acid in the yeast *bc<sub>1</sub>* complex [26]. Thirdly, lipids also mediate the division between a bipartite intra-membrane structure of the *b<sub>6</sub>f* complex, in which each monomer of the dimer consists of: (i) the conserved core consisting of the cyt *b* and subunit IV polypeptides, with four and three TMHs, respectively; (ii) the peripheral domain containing the single TMH of cyt *f* and the domain-swapped TMH of the ISP and the four small peripheral hydrophobic peptides, Pet G, L, M and N, the latter a set of single closely packed TMHs that define a unique ‘picket fence’ around each monomer. The spatial distribution of the subunit polypeptides and their B-factors are consistent with the concept of a two-tiered structure: (a) the polytopic core of the cyt *b<sub>6</sub>* and subunit IV subunits with four and three TMHs, respectively, the only polytopic subunits in the complex; and (b) a peripheral domain containing the single TMH of cyt *f*, ISP, Pet G, L, M and N. The function of the lipids in mediating interactions between the core and peripheral domain, and thus stabilizing the complex, is a consequence of the binding sites of the p-side lipids residing on residues from both the cyt *b<sub>6</sub>* and ‘subunit IV’ subunits of the core, and on the four single

TMH subunits, Pet G, L, M and N, in the total set of six single TMH subunits in the peripheral domain [18,25].

### (a) Two unique lipid functions

#### (i) Formation of a super-complex of the *b<sub>6</sub>f* complex with the photosystem I reaction centre

A curious situation exists for the function of the lipid-like β-carotene molecule. Carotenoids serve a universal role in photosynthesis as a quencher of the deleterious chlorophyll triplet state whose decay to the ground state can be coupled to the generation of excited state singlet oxygen, which is deleterious to plant and human tissue. Thus, on the one hand, the *b<sub>6</sub>f* complex contains a β-carotene whose function is ostensibly to carry out a triplet–triplet transfer and thus to quench the chlorophyll triplet state and thereby prevent the formation of singlet oxygen. However, crystal structures of the *b<sub>6</sub>f* complex from both the cyanobacteria and from a green alga are in agreement with a distance of closest approach of the single chlorophyll *a* in the structure and the one β-carotene of 14 Å [1,6,28]. Theoretically, this separation is much too large for efficient triplet–triplet transfer [29]. However, a significant extent of such energy transfer, measured on a rapid time scale, less than 8 ns, has been determined [28]. It is noted, however, that if the only function of the β-carotene would be to carry out triplet–triplet transfer, it should be in a position much closer to the chlorophyll *a* in the structure. Therefore, it is inferred that it has at least one additional function in the complex.

Considering its position in the structure, in which the β-carotene protrudes through the ‘picket fence’, between the Pet G and Pet M single TMHs by approximately 11 Å (figure 2), it is suggested that it could function as a ‘latch’ that connects to another structure in the photosynthetic membrane.




Figure 2. The exposed (11 Å) chain of  $\beta$ -carotene that is proposed to be a ‘latch’ to photosystem I, mediating the formation of a super-complex [30]. The  $b_6f$  complex has a peripheral  $\beta$ -carotene inserted into the trans membrane core of the complex (PDB 2E74). This pigment overlaps with the position of a peripheral cardiolipin of the yeast respiratory  $bc_1$  complex (PDB 3CX5) that mediates the formation of a  $bc_1$ –cyt  $c$  oxidase super-complex. The extension of the  $b_6f$   $\beta$ -carotene into the lipid bilayers is proposed to allow it to act as a structural antenna for the formation of  $b_6f$ –photosystem I super-complex. 2Fo–Fc map, 1.0 $\sigma$  (approx. 0.1 e<sup>−</sup>/Å<sup>3</sup>).

Examination of superimposed  $b_6f$  and  $bc_1$  structures indicates that the enigmatic  $\beta$ -carotene in the  $b_6f$  complex is in a position similar to that of a cardiolipin [31], which is positioned at the interface of a super-complex of the  $bc_1$  complex formed with cyt oxidase in the bovine respiratory chain [32,33]. It had previously been suggested that formation of a cyt  $b_6f$ –photosystem I (PSI) reaction centre ‘super-complex’ could be mediated by the  $\beta$ -carotene [6,34]. The  $b_6f$  complex in the green alga *C. reinhardtii* has been isolated in a large (‘super’) complex that includes the PSI reaction centre [30]. Thus, it is proposed that the surface-exposed and significantly protruding  $\beta$ -carotene in the peripheral domain of the  $b_6f$  complex is involved in mediating the formation of a super-complex with the PSI reaction centre complex.

(ii) *Lipid replacement of a trans membrane helix: trans-membrane signalling in the  $b_6f$  complex; an unusual transition in evolution*

It has long been known [35,36] that the eight TMHs of the cyt  $b$  subunit of the cyt  $bc_1$  complex are split into two subunits in the  $b_6f$  complex: firstly, the four TMH cyt  $b_6$  subunit, which corresponds to the N-terminal haem-binding four TMHs of the  $bc_1$  complex, and secondly subunit IV, consisting of three TMHs, the fourth largest subunit in the  $b_6f$  complex. Thus, the eight TMHs (A–H) in the cyt  $b$  subunit of the cyt  $bc_1$  complex are replaced by seven TMHs (A–G) in the  $b_6f$  complex. However, with no material in this niche formed by the eighth TMH in  $bc_1$ , a vacuum would result at this position in the structure

of the  $b_6f$  complex. Presumably, membrane protein structures abhor a vacuum (see discussion in [37]). It has recently been found that this space in the  $b_6f$  complex can largely be filled by a lipid and the bound chlorophyll  $a$  [25] (figure 3). The eight versus seven TMH problem in the cyt  $b$  polypeptide in energy-transducing membranes has previously been discussed in the evolutionary context of the residence of the split (four + three) seven TMHs of  $b_6f$ -like complexes in the ‘green clade’ [38]. In addition to the questions posed in [38] in the context of the finding that a lipid molecule in the  $b_6f$  complex occupies much of the space filled by the eighth TMH in the  $bc_1$  complex, one may ask: (i) which cyt  $bc$  complex appeared first in evolution, the respiratory or photosynthetic cyt  $bc$  complex? (ii) Thus, was the eighth TMH (the ‘H’ helix) replaced by two subunits consisting of seven TMHs, with a space in the structure for a lipid or another TMH from another subunit? (iii) Alternatively, was the sequence of events *vice versa*, i.e. the two subunits with seven TMHs replaced by a single cyt  $b$  subunit with eight TMHs? (iv) What was the ‘driving-force’ or ‘evolutionary pressure for one direction of the change or the other? A hypothesis for the answer to question (iv) is provided below.

Briefly summarizing available dating information and directions in the relevant branches of the evolutionary tree [38,39]: (i) carbon signatures implying established life in the absence of evidence for the presence of molecular oxygen have been found in Greenland rocks that have been dated to 3.8 Ga [40], implying the presence of bacteria containing chemo-lithotrophic respiratory chains; (ii) the presence of these bacteria substantially precedes that of the molecular oxygen (2.8–2.4 Ga [40]) that is associated with the existence of primitive oxygenic photosynthetic bacteria (i.e. cyanobacteria); (iii) the subunit structure of the  $b_6f$  complex from the cyanobacteria, with eight subunits, five redox prosthetic groups and seven prosthetic groups in total, has a greater degree of complexity than that of the proteobacterial  $bc_1$  complex that possesses three protein subunits and a total of four prosthetic groups [41]. Therefore, although not certain, it does not seem unlikely that the transition of the cyt  $bc$  complex in evolution was from the proto- $bc_1$  complex with an eight TMH cyt  $b$  polypeptide to the seven TMH  $b_6f$  complex present in the ‘green clade’, consistent with a ‘respiration-early’ hypothesis [42].

(b) *The evolutionary force in the cytochrome b transition: a hypothesis*

It is proposed that an evolutionary selection for this transition is trans membrane signalling in the  $b_6f$  complex in which the level of plastoquinol or quinol oxidation by the Rieske protein on the p-side of the complex can activate an enzyme bound on the n-side of the complex, studied mostly for the light harvesting complex II (LHCII) kinase [43–45] that regulates state transitions of light-harvesting chlorophyll proteins. The p-side site of the quinol oxidation by the [2Fe–2S] complex is proximal to the trans membrane F and G  $\alpha$ -helices, as seen in the structure of the  $b_6f$




Figure 3. The missing helix; trans membrane signalling. Substitution of the eighth helix in the  $bc_1$  complex with a lipid in  $b_6f$ . Trans membrane signalling by  $cyt\ b_6f$ ; p-side niche between the 'F' and 'C' TMHs defines a portal for passage of lipophilic quinol/quinone. (a) The eight TMHs of the respiratory  $bc_1$  complex and the seven (four + three) TMHs in  $b_6f$ ; (b) Domain containing F, G and H TMHs in the yeast  $bc_1$  complex and the F and G TMHs in  $b_6f$ . (b) binding niche of quinone analogue inhibitor, tridecyl-stigmatellin. (c) Ribbon diagram showing superposition of  $bc_1$  H-helix (orange) and, in  $b_6f$ , DOPC lipid (yellow) and the chlorophyll *a* chlorin ring and phytol tail (green). Binding site of p-side quinone analogue inhibitor, tridecyl-stigmatellin (blue). (d) Inferred structure–function equivalence of  $bc_1$  TMH 'H' and (lipid together with chlorophyll *a*) in  $b_6f$  complex.

complex. The niche unoccupied by the H-helix that is lost from the  $bc_1$  complex also spans the complex and the membrane at this site. This niche could be occupied by a trans membrane helical domain documented for the LHCII kinase (*stt7*) of *C. reinhardtii* (J.-D. Rochaix 2012, personal communication). However, along with four other proteins that have been found in isolated  $b_6f$  complex [7] before crystallization, the *stt7* LHCII protein kinase is not present in the crystals of the complex. It is presumed that such proteins are peripherally or loosely bound, which should not be the case for a subunit with a TMH. Alternatively, conformational changes associated with trans membrane signalling could be mediated by the intercalated lipid observed in the structure of the  $b_6f$  complex (PDB 2E74 [4,25]), perhaps coupled to the conformational flexibility of the chlorophyll *a* phytol chain that is inferred from the different positions of this chain in the structures of the complex from the cyanobacteria (PDB 2E74) and *C. reinhardtii* (PDB 1Q90).

A caveat for the inference of a first appearance of an LHC kinase in cyanobacteria during the course of evolution is that cyanobacteria do not display the state transitions characteristic of algae and higher plants that are associated with the function of the light-harvesting chlorophyll proteins in regulating the distribution of light energy to the two photosystems. However, a functionally similar regulatory mechanism

has been proposed for cyanobacteria [46,47], in which phosphorylation of the phycobiliproteins has been implicated [46]. Other redox-linked functions that have been proposed to be coupled to the redox state of the plastoquinone pool, and thereby quinol oxidation by the [2Fe–2S] Rieske protein, are chloroplast [48] and nuclear [49] gene expression, and other sensing functions [49].

These studies were supported by National Institutes of Health grant no. GM-038323. We thank J. Allen, R. E. Blankenship, and J.-D. Rochaix for helpful discussions.

## REFERENCES

- 1 Kurisu, G., Zhang, H., Smith, J. L. & Cramer, W. A. 2003 Structure of the cytochrome  $b_6f$  complex of oxygenic photosynthesis: tuning the cavity. *Science* **302**, 1009–1014. (doi:10.1126/science.1090165)
- 2 Yan, J., Kurisu, G. & Cramer, W. A. 2006 Structure of the cytochrome  $b_6f$  complex: binding site and intraprotein transfer of the quinone analogue inhibitor 2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone. *Proc. Natl Acad. Sci. USA* **103**, 67–74. (doi:10.1073/pnas.0504909102)
- 3 Yan, J., Kurisu, G. & Cramer, W. A. 2006 Intraprotein transfer of the quinone analogue inhibitor 2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone in the cytochrome  $b_6f$  complex. *Proc. Natl Acad. Sci. USA* **103**, 69–74. (doi:10.1073/pnas.0504909102)

- 4 Yamashita, E., Zhang, H. & Cramer, W. A. 2007 Structure of the cytochrome *b<sub>6</sub>f* complex: quinone analogue inhibitors as ligands of heme *c<sub>n</sub>*. *J. Mol. Biol.* **370**, 39–52. (doi:10.1016/j.jmb.2007.04.011)
- 5 Baniulis, D., Yamashita, E., Whitelegge, J. P., Zatsman, A. I., Hendrich, M. P., Hasan, S. S., Ryan, C. M. & Cramer, W. A. 2009 Structure–function, stability, and chemical modification of the cyanobacterial cytochrome *b<sub>6</sub>f* complex from *Nostoc* sp. PCC 7120. *J. Biol. Chem.* **284**, 9861–9869. (doi:10.1074/jbc.M809196200)
- 6 Stroebel, D., Choquet, Y., Popot, J.-L. & Picot, D. 2003 An atypical haem in the cytochrome *b<sub>6</sub>f* complex. *Nature* **426**, 413–418. (doi:10.1038/nature02155)
- 7 Cramer, W. A., Yamashita, E. & Hasan, S. S. 2011 The Q cycle of cytochrome *bc* complexes: a structure perspective. *Biochim. Biophys. Acta/Bioenergetics* **1807**, 788–802. (doi:10.1016/j.bbabi.2011.02.006)
- 8 Soriano, G. M., Smith, J. L. & Cramer, W. A. 2001 Cytochrome *f*. In *Handbook of metalloproteins* (eds A. Messerschmidt, R. Huber, K. Wieghardt & T. Poulos), pp. 172–181. London, UK: Wiley.
- 9 Carrell, C. J., Zhang, H., Cramer, W. A. & Smith, J. L. 1997 Biological identity and diversity in photosynthesis and respiration: structure of the lumen-side domain of the chloroplast Rieske protein. *Structure* **5**, 1613–1625. (doi:10.1016/S0969-2126(97)00309-2)
- 10 Cramer, W. A., Zhang, H., Yan, J., Kurisu, G. & Smith, J. L. 2006 Trans-membrane traffic in the cytochrome *b<sub>6</sub>f* complex. *Annu. Rev. Biochem.* **75**, 769–790. (doi:10.1146/annurev.biochem.75.103004.142756)
- 11 Nussberger, S., Dorr, K., Wang, D. N. & Kuhlbrandt, W. 1993 Lipid–protein interactions in crystals of the plant light-harvesting complex. *J. Mol. Biol.* **234**, 347–356. (doi:10.1006/jmbi.1993.1591)
- 12 McAuley, K. E., Fyfe, P. K., Ridge, J. P., Isaacs, N. W., Cogdell, R. J. & Jones, M. R. 1999 Structural details of an interaction between cardiolipin and an integral membrane protein. *Proc. Natl Acad. Sci. USA* **96**, 14 706–14 711. (doi:10.1073/pnas.96.26.14706)
- 13 Camara-Artigas, A., Brune, D. & Allen, J. P. 2002 Interaction between lipids and bacterial reaction centers determined by protein crystallography. *Proc. Natl Acad. Sci. USA* **99**, 11 055–11 060. (doi:10.1073/pnas.162368399)
- 14 Fyfe, P. K., Hughes, A. V., Heathcote, P. & Jones, M. R. 2005 Proteins, chlorophylls and lipids: X-ray analysis of a three-way relationship. *Trends Plant Sci.* **10**, 275–282. (doi:10.1016/j.tplants.2005.04.007)
- 15 Jones, M. R. 2007 Lipids in photosynthetic reaction centres: structural roles and functional holes. *Prog. Lipid Res.* **46**, 56–87. (doi:10.1016/j.plipres.2006.06.001)
- 16 Wöhri, A. B., Wahlgren, W. Y., Malmerberg, E., Johansson, L. C., Neutze, R. & Katona, G. 2009 Lipidic sponge phase crystal structure of a photosynthetic reaction center reveals lipids on the protein surface. *Biochemistry* **48**, 9831–9838. (doi:10.1021/bi900545e)
- 17 Guskov, A., Kern, J., Gabdulkhakov, A., Broser, M., Zouni, A. & Saenger, W. 2009 Cyanobacterial photosystem II at 2.9-Å resolution and the role of quinones, lipids, channels and chloride. *Nat. Struct. Mol. Biol.* **16**, 334–342. (doi:10.1038/nsmb.1559)
- 18 Kern, J., Zouni, A., Guskov, A. & Krauss, N. 2009 Lipids in the structure of the photosystem I, photosystem II, and the cytochrome *b<sub>6</sub>f* complex. In *Lipids in photosynthesis: essential and regulatory functions* (ed. H. W. a. N. Murata), pp. 203–242. Berlin, Germany: Springer Science.
- 19 Umena, Y., Kawakami, K., Shen, J. R. & Kamiya, N. 2011 Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 Å. *Nature* **473**, 55–60. (doi:10.1038/nature09913)
- 20 Jordan, P., Fromme, P., Witt, H. T., Klukas, O., Saenger, W. & Krauss, N. 2001 Three-dimensional structure of cyanobacterial photosystem I at 2.5 Å resolution. *Nature* **411**, 909–917. (doi:10.1038/35082000)
- 21 Shinzawa-Itoh, K. *et al.* 2007 Structures and physiological roles of 13 integral lipids of bovine heart cytochrome *c* oxidase. *EMBO J.* **26**, 1713–1725. (doi:10.1038/sj.emboj.7601618)
- 22 Qin, L., Hiser, C., Mulichak, A., Garavito, R. M. & Ferguson-Miller, S. 2006 Identification of conserved lipid/detergent binding site in a high resolution structure of the membrane protein, cytochrome *c* oxidase. *Proc. Natl Acad. Sci. USA* **103**, 16 117–16 122. (doi:10.1073/pnas.0606149103)
- 23 Qin, L., Sharpe, M. A., Garavito, R. M. & Ferguson-Miller, S. 2007 Conserved lipid-binding sites in membrane proteins: a focus on cytochrome *c* oxidase. *Curr. Opin. Struct. Biol.* **17**, 444–450. (doi:10.1016/j.sbi.2007.07.001)
- 24 Palsdottir, H. & Hunte, C. 2004 Lipids in membrane protein structures. *Biochim. Biophys. Acta* **1666**, 2–18. (doi:10.1016/j.bbamem.2004.06.012)
- 25 Hasan, S. S., Yamashita, E., Ryan, C. M., Whitelegge, J. P. & Cramer, W. A. 2011 Conservation of lipid functions in cytochrome *bc* complexes. *J. Mol. Biol.* **414**, 145–162. (doi:10.1016/j.jmb.2011.09.023)
- 26 Solmaz, S. R. & Hunte, C. 2008 Structure of complex III with bound cytochrome *c* in reduced state and definition of a minimal core interface for electron transfer. *J. Biol. Chem.* **283**, 17 542–17 549. (doi:10.1074/jbc.M710126200)
- 27 de Vitry, C., Ouyang, Y., Finazzi, G., Wollman, F. A. & Kallas, T. 2004 The chloroplast Rieske iron–sulfur protein. At the crossroad of electron transport and signal transduction. *J. Biol. Chem.* **279**, 44 621–44 627. (doi:10.1074/jbc.M406955200)
- 28 Kim, H., Dashdorj, N., Zhang, H., Yan, J., Cramer, W. A. & Savikhin, S. 2005 An anomalous distance dependence of intra-protein chlorophyll–carotenoid triplet energy transfer. *Biophys. J.* **89**, 28–30. (doi:10.1529/biophysj.105.069609)
- 29 Dexter, D. L. 1953 A theory of sensitized luminescence in solids. *J. Chem. Phys.* **21**, 836–850. (doi:10.1063/1.1699044)
- 30 Iwai, M., Takizawa, K., Tokutsu, R., Okamuro, A., Takahashi, Y. & Minagawa, J. 2010 Isolation of the supercomplex that drives cyclic electron flow in photosynthesis. *Nature* **464**, 1210–1213. (doi:10.1038/nature08885)
- 31 Wenz, T., Hielscher, R., Hellwig, P., Schagger, H., Richters, S. & Hunte, C. 2009 Role of phospholipids in respiratory cytochrome *bc<sub>1</sub>* complex catalysis and supercomplex formation. *Biochim. Biophys. Acta* **1787**, 609–616. (doi:10.1016/j.bbabi.2009.02.012)
- 32 Schafer, E., Dencher, N. A., Vonck, J. & Parcej, D. N. 2007 Three-dimensional structure of the respiratory chain supercomplex I1III2IV1 from bovine heart mitochondria. *Biochemistry* **46**, 12 579–12 585. (doi:10.1021/bi700983h)
- 33 Althoff, T., Mills, D. J., Popot, J. L. & Kuhlbrandt, W. 2011 Arrangement of electron transport chain components in bovine mitochondrial supercomplex I1III2IV1. *EMBO J.* **30**, 4652–4664. (doi:10.1038/emboj.2011.324)
- 34 Kuhlbrandt, W. 2003 Structural biology: dual approach to a light problem. *Nature* **426**, 399–400. (doi:10.1038/426399a)
- 35 Widger, W. R., Cramer, W. A., Herrmann, R. G. & Trebst, A. 1984 Sequence homology and structural similarity between the *b* cytochrome of mitochondrial complex III and the chloroplast *b<sub>6</sub>f* complex: position of the

- cytochrome *b* hemes in the membrane. *Proc. Natl Acad. Sci. USA* **81**, 674–678. (doi:10.1073/pnas.81.3.674)
- 36 Schutz, M. *et al.* 2000 Early evolution of cytochrome *bc* complexes. *J. Mol. Biol.* **300**, 663–675. (doi:10.1006/jmbi.2000.3915)
- 37 Fyfe, P. K., Potter, J. A., Cheng, J., Williams, C. M., Watson, A. J. & Jones, M. R. 2007 Structural responses to cavity-creating mutations in an integral membrane protein. *Biochemistry* **46**, 10 461–10 472. (doi:10.1021/bi701085w)
- 38 Nitschke, W., van Lis, R., Schoepp-Cothenet, B. & Baymann, F. 2010 The ‘green’ phylogenetic clade of Rieske/cyt *b* complexes. *Photosyn. Res.* **104**, 347–355. (doi:10.1007/s11120-010-9532-1)
- 39 Hohmann-Marriott, M. F. & Blankenship, R. E. 2011 Evolution of photosynthesis. *Annu. Rev. Plant Biol.* **62**, 515–548. (doi:10.1146/annurev-arplant-042110-103811)
- 40 Blankenship, R. E. *et al.* 2011 Comparing photosynthetic and photovoltaic efficiencies and recognizing the potential for improvement. *Science* **332**, 805–809. (doi:10.1126/science.1200165)
- 41 Esser, L., Elberry, M., Zhou, F., Yu, C. A., Yu, L. & Xia, D. 2008 Inhibitor-complexed structures of the cytochrome *bc<sub>1</sub>* from the photosynthetic bacterium *Rhodobacter sphaeroides*. *J. Biol. Chem.* **283**, 2846–2857. (doi:10.1074/jbc.M708608200)
- 42 Castresana, J. & Saraste, M. 1995 Evolution of energetic metabolism: the respiration-early hypothesis. *Trends Biochem. Sci.* **20**, 443–448. (doi:10.1016/S0968-0004(00) 89098-2)
- 43 Vener, A. V., van Kan, P. J., Rich, P. R., Ohad, I. & Andersson, B. 1997 Plastoquinol at the quinol oxidation site of reduced cytochrome *bf* mediates signal transduction between light and protein phosphorylation: thylakoid protein kinase deactivation by a single-turnover flash. *Proc. Natl Acad. Sci. USA* **94**, 1585–1590. (doi:10.1073/pnas.94.4.1585)
- 44 Zito, F., Finazzi, G., Delosme, R., Nitschke, W., Picot, D. & Wollman, F. A. 1999 The Qo site of cytochrome *b<sub>6</sub>f* complexes controls the activation of the LHCII kinase. *EMBO J.* **18**, 2961–2969. (doi:10.1093/emboj/18.11.2961)
- 45 Zer, H. & Ohad, I. 2003 Light, redox state, thylakoid-protein phosphorylation and signaling gene expression. *Trends Biochem. Sci.* **28**, 467–470. (doi:10.1016/S0968-0004(03)00173-7)
- 46 Mullineaux, C. W. & Emllyn-Jones, D. 2005 State transitions: an example of acclimation to low-light stress. *J. Exp. Bot.* **56**, 389–393. (doi:10.1093/jxb/eri064)
- 47 Bernat, G., Schreiber, U., Sendtko, E., Stadnichuk, I. N., Rexroth, S., Rogner, M. & Koenig, F. 2012 Unique properties vs. common themes: the atypical cyanobacterium *Gloeobacter violaceus* PCC 7421 is capable of state transitions and blue-light-induced fluorescence quenching. *Plant Cell Physiol.* **53**, 528–542. (doi:10.1093/pcp/pcs009)
- 48 Allen, J. F., de Paula, W. B., Puthiyaveetil, S. & Nield, J. 2011 A structural phylogenetic map for chloroplast photosynthesis. *Trends Plant Sci.* **16**, 645–655. (doi:10.1016/j.tplants.2011.10.004)
- 49 Rochaix, J. D. 2011 Regulation of photosynthetic electron transport. *Biochim. Biophys. Acta* **1807**, 375–383. (doi:10.1016/j.bbabi.2010.11.010)